Fisheries Research, 17 (1993) 387–389
Elsevier Science Publishers B.V.

Contents of *Fisheries Research*, Volume 17

VOL. 17 NOS. 1–2 **JUNE 1993**

Special Issue: Pathological Conditions of Wild Salmonids

Introduction
Pathological conditions of wild salmonids
D.W. Bruno (Aberdeen, UK) 1

Ulcerative dermal necrosis (UDN) in wild salmonids
R.J. Roberts (Stirling, UK) 3
The history and current status of *Renibacterium salmoninarum*, the causative agent of bacterial kidney disease in Pacific salmon
J.L. Fryer (Corvallis, OR, USA) and C.N. Lannan (Newport, OR, USA) 15
Endocrine-induced pathology in stressed salmonid fish
A.D. Pickering (Ambleside, UK) 35
Proliferative kidney disease in wild salmonids
S.W. Feist and D. Bucke (Weymouth, UK) 51
Species of *Gyrodactylus* (Platyhelminthes, Monogenea) on salmonids in Sweden
G. Malmberg and M. Malmberg (Stockholm, Sweden) 59
Comparative susceptibility of native Scottish and Norwegian stocks of Atlantic salmon, *Salmo salar* L., to *Gyrodactylus salaris* Malmberg: Laboratory experiments
T.A. Bakke (Oslo, Norway) and K. MacKenzie (Aberdeen, UK) 69
Regulatory processes in the monogenean *Gyrodactylus salaris* Malmberg–Atlantic salmon (*Salmo salar* L.) association. I. Field studies in southeast Norway
P.A. Jansen and T.A. Bakke (Oslo, Norway) 87
Regulatory processes in the monogenean *Gyrodactylus salaris* Malmberg–Atlantic salmon (*Salmo salar* L.) association. II. Experimental studies
P.A. Jansen and T.A. Bakke (Oslo, Norway) 103
Piscirickettsia salmonis, a major pathogen of salmonid fish in Chile
C.N. Lannan (Newport, OR, USA) and J.L. Fryer (Corvallis, OR, USA) 115
The pattern of natural mortality throughout the life cycle in contrasting populations of brown trout, *Salmo trutta* L.
J.M. Elliott (Ambleside, UK) 123
An epidemiological study of furunculosis in salmon propagation in Japanese rivers
T. Nomura (Sapporo, Japan), M. Yoshimizu and T. Kimura (Hakodate, Japan) 137
Systemic, granulomatous hexamitosis of farmed Atlantic salmon: Interaction with wild fish
T.T. Poppe and T.A. Mo (Oslo, Norway) 147
VHS in wild living fish and experimental transmission of the virus
P.-J. Enzmann, M. Konrad and K. Parey (Tübingen, Germany) 153
Surveillance and control of infectious hematopoietic necrosis virus (IHNV) and

Oncorhynchus masou virus (OMV) of wild salmonid fish returning to the northern part of Japan 1976–1991
M. Yoshimizu, T. Nomura, Y. Ezura and T. Kimura (Hokkaido, Japan) 163
Diseases of wild sea trout in Scotland in relation to fish population decline
A.H. McVicar, L.A. Sharp (Aberdeen, UK), A.F. Walker (Pitlochry, UK) and A. W. Pike (Aberdeen, UK) 175
Production of nauplii of *Lepeophtheirus salmonis* (Krøyer) (Copepoda: Caligidae) from farmed and wild salmon and its relation to the infestation of wild sea trout (*Salmo trutta* L.) off the west coast of Ireland in 1991
O. Tully (Dublin, Ireland) and K.F. Whelan (Newport, Ireland) 187
Furunculosis as a possible factor in the decline of sea trout in the West of Ireland
J.F. McArdle, C. Dooley-Martyn, F. Geoghegan, F. McKiernan (Dublin, Ireland) and H. Rodger (Galway, Ireland) 201
The significance of diseases and anomalies in wild salmonids
D. Bucke (Weymouth, UK) 209
Parasites of Norwegian freshwater salmonids and interactions with farmed salmon—a review
G.A. Bristow (Bergen, Norway) 219
EC Directive 91/67/EEC and its relevance to disease in wild salmonids
E.B. Hudson and B.J. Hill (Weymouth, UK) 229
Modelling the impact of disease-induced mortality on the population size of wild salmonids
S. des Clers (London, UK) 237

Announcement from the Publisher 249

VOL. 17 NOS. 3–4 AUGUST 1993

Optic critical duration and contrast thresholds in the freshwater fish, *Lepomis macrochirus*, as determined behaviourally
G. Kawamura (Shimoarata, Kagoshima, Japan) and T. Shimowada (Tokyo, Japan) 251
Norway lobster (*Nephrops norvegicus*) from the southwest and south of Portugal — estimation of the effects of changing trawl mesh size and fishing effort by length cohort analysis
F. Cardador (Lisbon, Portugal) 259
The derivation of and changes in *Nephrops* per unit effort values for the Northern Ireland fishing fleet
D.A. Stewart (Belfast, UK), D. Agnew (Hobart, Tas., Australia), R. Boyd, R. Briggs and P. Toland (Belfast, UK) 273
A method to estimate the S–R relationship using rough information on the rate of exploitation
X. Li (Sendai, Japan) and P. Safran (Paris, France) 293
A comparison of one-tenth and full-scale measurements of the drag and geometry of a pelagic trawl
J.N. Ward (Hull, UK) and R.S.T. Ferro (Aberdeen, UK) 311
An analysis based on trawl-survey data of the state of the 'Italian' stock of *Mullus barbatus* in the Sicilian Channel, including management advice
D. Levi (Mazara del Vallo, Italy), M.G. Andreoli (Palermo, Italy) and G.B. Giusto (Mazara del Vallo, Italy) 333
Selection of the appropriate duration of experiments to measure the survival of animals

discarded from trawlers
T.J. Wassenberg and B.J. Hill (Cleveland, Qld., Australia) 343
Migration and growth of two tropical penaeid shrimps within Torres Strait, northern Australia
R.A. Watson and C.T. Turnbull (Cairns, Qld., Australia) 353

Short Communications
Observations on the distribution of lumpsucker (*Cyclopterus lumpus*, L.) in the Norwegian Sea
J.C. Holst (Bergen, Norway) 369
Analysis of trawl selectivity studies (addendum): implementation in SAS
R.B. Millar (St. John's, Nfld., Canada) 373

Book Reviews
Genetics and fish breeding 379
Introductions of exotic species 380

Announcement from the Publisher 383

Author Index 384